

# DUSTMARKS AND OZONE TATTOOS: AUTOGRAPHIC DISPLAYS OF AIR POLLUTION

**Dietmar Offenhuber**  
Northeastern University  
360 Huntington Ave.  
Boston MA 02139  
d.offenhuber@northeastern.edu

## **Abstract**

We present two examples of autographic visualizations - displays based on material traces rather than digital data - that aim to make environmental pollution visible and legible. Using particulate matter and ground-level ozone pollution as case studies, this pictorial illustrates the design principles of creating autographic visualizations that make phenomena reveal themselves.

## **Authors Keywords**

Autographic visualization, indexical visualization, physicalization, critical data studies, environmental phenomena

## **Introduction**

The public art projects *Staubmarke* and *Ozone Tattoo* are autographic displays that make aspects of air pollution—usually represented in abstract metrics—accessible to sensory experience. In both projects, the respective phenomena are made to reveal themselves in material traces, unmediated by symbolically encoded data.




Focusing on dust and particulate matter, *Staubmarke* (figure above) makes urban air pollution visible by introducing a visual system to observe and compare the dust patina across urban surfaces and contrast it with data from a citizen science sensor network.



*Ozone Tattoo* (figure above) uses plants as sensors for ground-level ozone pollution. Similar to *Staubmarke*, a visual reference pattern shows how ozone damage manifests on the leaves of certain indicator plants (tobacco, snap bean), helping to detect ozone episodes through comparison with the untreated leaves.

## Data Visualization vs. Autographic Visualization



**Data visualization** is confined within the borders of symbolic representation. It can only start when data already exist. Therefore, the circumstances of data generation and the precise relationship of a data set with the phenomenon it describes often remain obscure. While data sets often contain signatures that point to how they were collected, a gap remains between data and the phenomenon of interest. Because of this gap, we often find it difficult to reconcile the results of a data analysis with experiential phenomena, especially in the cases of environmental pollution or global warming.

**Autographic (and indexical) visualization** operate at the material bases of data: physical traces and other forms of environmental information [7,8,9]. Autographic visualization is defined as “a set of techniques for revealing material phenomena as visible traces and guiding their interpretation. Designing an autographic display means setting the conditions that allow a trace to emerge” [7]. A trace is understood as any material structure or constellation that can present itself to the senses. While data visualization supports the interpretation of data, autographic visualization makes the origin of a datum and the process of its generation legible. Since data analysis is more versatile than the manipulation of physical traces, a gap remains between data and analysis.

**A combined model** of symbolic data analysis and visualization with an autographic display of the underlying physical phenomena allows a transgression of the boundary between the world of symbols and the physical realm. It offers a critical approach that makes visible the assumptions and decisions behind acts of measurement and calls attention to the material nature and contexts of data.


Figure: Conceptual model of InfoVis, autographic visualization and a hybrid model, from [7]

**Staubmarke (dustmark)**

<http://dust.zone>

Public space installation

Drehmoment Festival Stuttgart, October 2018






Stuttgart is one of Germany's cities with the worst air pollution, suffering under frequent episodes of particulate matter pollution exceeding the WHO criteria values.

Since the city is a major center of the German car industry, blamed by many for the environmental condition, air quality is an intensely political issue in Stuttgart.


In the course of this controversy, citizens often mistrust and critique official measurements and sensing methodologies. In response, citizen science enthusiasts have founded *luftdaten.info*, a sensing initiative of self-assembled air-quality sensors and an online platform for archiving and disseminating crowd-sourced airquality data.



It is striking that such public controversies often quickly converge on questions of the appropriate methods and thresholds of measurement [4]. It does not help that common metrics for measuring particulate matter are highly reductive, representing only certain qualities of the invisible phenomenon. These circumstances make it tempting to treat air pollution as a single, abstract number.

Above: community-built dust sensor by @airrohr, source: <https://twitter.com/airrohr/status/914154105941237760>

Below: staubmarke website with live data from <http://luftdaten.info>



At the same time, particulate matter is a physical material, visible on used dustmasks, and accumulating on all surfaces of the city.

Staubmarke introduces a uniform visual pattern to make the urban patina and the process of its accumulation legible. Using the technique of reverse graffiti, which creates visible patterns by partially cleaning surfaces rather than applying paint, visual markers are applied in public spaces of the city.

The chosen locations are monitored by airquality sensors from the *luftdaten.info* initiative, and the highest daily average value during the past month is displayed on the graffiti.

As more dust accumulates, the dustmarks will slowly fade away. Due to the design of the pattern, the finest parts will disappear first, allowing to estimate the speed of the process. Observation of the patterns over the following year will offer new insight into this process.

It might be objected that the patina consists of more than just particulate matter. This is true, but also official metrics do not capture the phenomenon in its complexity. Staubmarke is meant to inspire reflection about the materiality of data and their processes of measurement.

Top-left: sample of particulate matter for gravimetric analysis, courtesy Frank M. Rauch, wikimedia.

All other photos by the author.

# Dustmarks, four months after their application

Fotos: Frank Kleinbach




## Anatomy of a dustmark



**Reference pattern**  
Making the degree of dust accumulation visible, legible and comparable.

**Gradient**  
The finest part of the pattern will disappear first, revealing the process of reaccumulation



**Contextualizing digital information**  
Link to live visualization, and sensor value.

## Ozone Tattoo

Public space installation  
Cambridge Arts Council, May 2019

Project partners:  
Dr. Vehram Elagoz, Lesley University  
Dr. Kent Burkey, USDA ARS Maryland








Ozone fumigator to create reference damage patterns

Groundlevel ozone a serious pollutant that puts humans, plants, and animals at risk. It is a side effect of air pollution associated with fossil fuels and is connected to global warming. Groundlevel ozone emerges mainly on hot summer days. Researchers estimate that the increasing intensity of summer heat waves will lead to a 70-100% increase in ozone episodes [10].

Compared to particulate matter, ground-level ozone is more difficult to measure and rarely covered by air quality stations. Ozone ( $O_3$ ) is highly volatile, and calibration issues make sensors unaffordable for amateurs.



Ozone episodes can be reliably detected by observing certain indicator plants, including tobacco, snap beans, or cone flowers. Each of them responds with characteristic damage patterns.

*Ozone Tattoo* uses plants as bio-indicators to visualize ozone pollution. It is based on the established citizen science approach of planting ozone gardens, which allow communities to monitor pollution by observing plants that are sensitive to ozone and react with visible symptoms [3]. To make the damage to plants more legible, we developed the novel method of ozone tattoos. The tattoos are damage patterns on the surface of the leaves, created by localized exposure to ozone. These damage patterns act as a biological visualization that allows us to decode the impact of the pollutant through visual comparison with the healthy portions of the leaf.



Reference damage pattern for snap beans

Healthy leaves for comparison (will show similar patterns after ozone episode)




The ozone tattoo machine uses a weak ozone generator based on electric corona discharge, a fan and a tubing system of teflon and silicone tubes that bring the ozone to the leaf before it dissipates. Final ozone concentrations used were between 0.1 and 0.3 ppm.

leaf areas exposed to ozone

Exposure to  
0.4 ppm  
Ozone for 24

Tobacco  
O<sub>3</sub> sensitive  
O<sub>3</sub> tolerant

Snap bean  
O<sub>3</sub> sensitive  
O<sub>3</sub> tolerant



Ozone indicator system based on a combination of ozone-sensitive and tolerant variations of the same plant. Comparison of ozone damage on both allows estimating the ozone concentration [1,2,5].

Seeds provided by USDA Agricultural Research Service (ARS):

- Snap beans: S156 (O<sub>3</sub> sensitive) and R123 (O<sub>3</sub> tolerant) genotypes of *Phaseolus vulgaris* L.
- Tobacco: Bel-W3 (O<sub>3</sub> sensitive) and Bel-B (O<sub>3</sub> tolerant)

**Ozone Tattoo on a tobacco leaf**

Tobacco BEL-W3 (O<sub>3</sub> - sensitive)



## Experimental Ozone Tattoos with different fumigator end pieces

Tobacco BEL-W3 ( $O_3$  sensitive)



This pattern indicates acute rather than chronic ozone damage, the damaged areas are larger due to higher ozone concentration.


## Ozone Tattoo resin casts

Documenting different ozone tattoo experiments as physical specimen.





Ozone tattoo machine and tobacco plants exhibited at Cambridge Art Council Gallery, <https://www.untoldpossibilities.org>



Views of leaf damage on snap bean and tobacco plants through a light microscope views.

## Ozone Tattoo Ozone Garden

Different sections of O<sub>3</sub> sensitive and tolerant varieties of tobacco and snap bean allow the detection of ozone episodes and the estimation of their severity. The tattoos on each plant help identifying the damage and serve as a contrast for comparison.



## References

1. Burkey, K.O. et al. 2012. Field assessment of a snap bean ozone bioindicator system under elevated ozone and carbon dioxide in a free air system. *Environmental Pollution*. 166, (Jul. 2012), 167-171.
2. Elagöz, V. and Manning, W.J. 2005. Responses of sensitive and tolerant bush beans (*Phaseolus vulgaris* L.) to ozone in open-top chambers are influenced by phenotypic differences, morphological characteristics, and the chamber environment. *Environmental Pollution*. 136, 3 (Aug. 2005), 371-383.
3. Fishman, J. et al. 2014. The St. Louis ozone gardens: Visualizing the impact of a changing atmosphere. *Bulletin of the American Meteorological Society*. 95, 8 (2014), 1171-1176.
4. Gabrys, J. 2016. Practicing, materialising and contesting environmental data. *Big Data & Society*. 3, 2 (Nov. 2016).
5. Heggstad, H.E. 1991. Origin of Bel-W3, Bel-C and Bel-B tobacco varieties and their use as indicators of ozone. *Environmental Pollution*. 74, 4 (Jan. 1991), 264-291.
6. Kuznetsov, S. et al. 2011. Nurturing natural sensors. *Proceedings of the 13th international conference on Ubiquitous computing (New York, NY, USA, 2011)*, 227-236.
7. Offenhuber, D. Forthcoming. "Data by Proxy – Material Traces as Autographic Visualizations," *IEEE Transactions on Visualization and Computer Graphics (Proceedings of InfoVis 2019)*.
8. Offenhuber, D. and Telhan, O. 2015. Indexical Visualization—the Data-less Information Display. U. Ekman et al., eds. Routledge. 288-303.
9. Schofield, T. et al. 2013. Indexicality and visualization: exploring analogies with art, cinema and photography. *Proceedings of the 9th ACM Conference on Creativity & Cognition (2013)*, 175-184.
10. Shen, L. et al. 2016. Impact of increasing heat waves on U.S. ozone episodes in the 2050s: Results from a multimodel analysis using extreme value theory. *Geophysical Research Letters*. 43, 8 (2016), 4017-4025.